

INVARIANT THEORY FOR QUANTUM WEYL ALGEBRAS UNDER FINITE GROUP ACTION

S. CEKEN, J. H. PALMIERI, Y.-H. WANG AND J. J. ZHANG

ABSTRACT. We study the invariant theory of a class of quantum Weyl algebras under group actions and prove that the fixed subrings are always Gorenstein. We also verify the Tits alternative for the automorphism groups of these quantum Weyl algebras.

INTRODUCTION

Fix a field k . For $n \geq 2$, let W_n be the (-1) -quantum Weyl algebra: this is the k -algebra generated by $x_1, \dots, x_n$ subject to the relations $x_i x_j + x_j x_i = 1$ for all $i \neq j$.

Theorem 1. *Assume that $\text{char } k = 0$. Let n be an even integer ≥ 4 and let G be a group acting on W_n . Then the fixed subring W_n^G under the G -action is filtered Artin-Schelter Gorenstein.*

The above theorem was announced in [CPWZ1, Theorem 2] without proof. The first purpose of this paper is to provide a proof of this, using some earlier results from noncommutative invariant theory [JZ, KKZ1]. Secondly, we discuss the automorphism group of W_n when $n \geq 3$ is odd [Theorem 2]. We also want to correct some small errors in [CPWZ1]: see Remarks 1.5(2) and 2.7. Finally, we give a criterion for an isomorphism question for a class of (-1) -quantum Weyl algebras [Theorem 3].

One aspect of invariant theory is to study homological properties of fixed subrings (also called invariant subrings) under group actions. When A is regular (or has finite global dimension), the fixed subring A^G under the action of a finite group G is Cohen-Macaulay when $\text{char } k$ does not divide the order of G . One interesting question is *when a fixed subring A^G is Gorenstein*. The famous Watanabe theorem [Wa] answers such a question for commutative polynomial rings. Theorem 1 above provides a solution for W_n when n is even. However, this question is open for W_n^G when n is odd (even when $n = 3$ and G is finite). Another interesting question is *when a fixed subring A^G is regular*. The classical result of Shephard-Todd-Chevalley [ST, Ch] answers this question for commutative polynomial rings. The question is open for W_n for all $n \geq 3$, though we have an easy partial result: see Proposition 1.8. A recent survey on invariant theory of Artin-Schelter regular algebras is given by Kirkman in [Ki].

Another aspect of invariant theory is to study the structure of the automorphism group $\text{Aut}(A)$ of an algebra A . There is a long history and an extensive

2000 *Mathematics Subject Classification.* Primary 16W20, 16E65.

Key words and phrases. automorphism group, quantum Weyl algebra, Artin-Schelter Gorenstein property, free subgroup.

study of the automorphism groups of algebras. Determining the full automorphism group of an algebra is generally a notoriously difficult problem. Recently, significant progress has been made in finding the full automorphism groups of some noncommutative algebras. For example, during the last two years, Yakimov has proved the Andruskiewitsch-Dumas conjecture and the Launois-Lenagan conjecture by using a rigidity theorem for quantum tori [Y1, Y2]. The automorphism groups of generalized or quantum Weyl algebras have been studied by several researchers [AD, BJ, SAV]. The authors used the discriminant method to determine the automorphism groups of some noncommutative algebras [CPWZ1, CPWZ2]. When n is even, $\text{Aut}(W_n)$ was worked out by the authors in [CPWZ1, Theorem 1]. Unfortunately, we have not been able to determine the full automorphism group of W_n when n is odd [CPWZ1, Example 5.10 and Question 5.13].

The second theorem concerns $\text{Aut}(W_n)$ when n is odd. We will consider a slightly more general setting. Let $\mathcal{A} := \{a_{ij} \in k \mid 1 \leq i < j \leq n\}$ be a set of scalars. Define a modified (-1) -quantum Weyl algebra $V_n(\mathcal{A})$ [CPWZ1, Section 4] to be the k -algebra generated by $\{x_1, \dots, x_n\}$ subject to the relations

$$x_i x_j + x_j x_i = a_{ij}, \quad \forall 1 \leq i < j \leq n.$$

As a special case, note that if $a_{ij} = 1$ for all $i < j$, then $V_n(\mathcal{A}) = W_n$. Another special case is when $a_{ij} = 0$ for all $i < j$, in which case $V_n(\mathcal{A})$ is just the skew polynomial ring $k_{-1}[x_1, \dots, x_n]$.

Theorem 2. *Let $n \geq 3$ be an odd integer. Suppose $\text{char } k$ does not divide $(n-1)!$. Then $\text{Aut}(V_n(\mathcal{A}))$ contains a free group on two generators. As a consequence, it contains a free group on countably many generators.*

Note that [CPWZ1, Theorem 1] implies that, when n is even, $\text{Aut}(W_n)$ is finite (and so virtually solvable). A more general statement for $\text{Aut}(V_n(\mathcal{A}))$ is Lemma 1.1. Combining Theorem 2 with Lemma 1.1(2,3), we obtain that in characteristic 0, $\text{Aut}(V_n(\mathcal{A}))$ either is virtually solvable or contains a free subgroup of rank two. This can be viewed as a version of the Tits alternative [Ti] for the automorphism groups of $V_n(\mathcal{A})$. The original Tits alternative states that every finitely generated linear group is either virtually solvable or contains a free subgroup of rank two. We also have a version of the Tits alternative for the class of the automorphism groups of skew polynomial rings in a separate paper [CPWZ3].

In Section 3, we use the discriminant to prove the following criterion for when two $V_n(\mathcal{A})$ s are isomorphic (in the case when n is even). For simplicity, let $a_{ij} = a_{ji}$ if $i > j$.

Theorem 3. *Suppose $\text{char } k \neq 2$ and let n be an even integer. Let $\mathcal{A}' := \{a'_{ij} \mid 1 \leq i < j \leq n\}$ be another set of scalars in k . Then $V_n(\mathcal{A}) \cong V_n(\mathcal{A}')$ if and only if there are a permutation $\sigma \in S_n$ and nonzero scalars λ_i for $i = 1, \dots, n$ such that $a'_{ij} = \lambda_i \lambda_j a_{\sigma(i)\sigma(j)}$ for all i and j .*

As a consequence of Theorem 3, when $n \geq 4$ is even, there are infinitely many non-isomorphic $V_n(\mathcal{A})$ s.

In Section 4, we give some examples of $\text{Aut}(V_n(\mathcal{A}))$ for $n = 4, 6$ and list some questions about $V_n(\mathcal{A})$.

1. PROOF OF THEOREM 1

Throughout let k be a base commutative domain. Modules, vector spaces, algebras, and morphisms are over k . In this section we further assume that k is a field.

As above, for each $n \geq 2$, let $V_n(\mathcal{A})$ denote the algebra generated by $\{x_i\}_{i=1}^n$ subject to the relations $x_i x_j + x_j x_i = a_{ij}$ for all $i \neq j$. This is a filtered Artin-Schelter regular algebra in the following sense. We refer to [KKZ2, Definition 1.2] for the definition of (connected graded) Artin-Schelter regularity. Let $F := \{F_n \subseteq A \mid n \geq 0\}$ be an increasing filtration on an algebra A satisfying

- (a) $F_0 = k$,
- (b) $F_n F_m \subseteq F_{n+m}$ for all $n, m \geq 0$,
- (c) $A = \bigcup_n F_n$.

The associated graded algebra with respect to F , denoted by $\text{gr}_F A$, is defined to be $\text{gr}_F A = \bigoplus_{i=0}^{\infty} F_i / F_{i-1}$. This is a connected graded algebra by condition (a). An algebra A is called *filtered Artin-Schelter regular* (resp. *filtered Artin-Schelter Gorenstein*) if there is a filtration F such that $\text{gr}_F A$ is Artin-Schelter regular (resp. Artin-Schelter Gorenstein). In our case, we filter $V_n(\mathcal{A})$ by setting $F_0 = k$, $F_1 = k + \sum_{s=1}^n k x_s$, and $F_i = (F_1)^i$ for all $i \geq 2$. Then it is easy to see that $\text{gr}_F V_n(\mathcal{A}) \cong k_{-1}[x_1, \dots, x_n] (= V_n(\{0\}))$, the skew polynomial ring generated by $\{x_i\}_{i=1}^n$ subject to the relations $x_j x_i = -x_i x_j$ for all $i < j$. Since $k_{-1}[x_1, \dots, x_n]$ is Artin-Schelter regular, $V_n(\mathcal{A})$ is filtered Artin-Schelter regular. The symmetric group S_n acts on the set $\{x_i\}_{i=1}^n$ naturally. It is easy to see that this action extends to an S_n -action on the algebras $V_n(\mathcal{A})$ and $k_{-1}[x_1, \dots, x_n]$ uniquely; note that the invariant theory of S_n -actions on $k_{-1}[x_1, \dots, x_n]$ was studied in [KKZ1].

The group of all affine automorphisms of $V_n(\mathcal{A})$, denoted by $\text{Aut}_{\text{af}}(V_n(\mathcal{A}))$, can be worked out by using easy combinatorics [CPWZ1, Lemma 4.3]. Let $G(\mathcal{A})$ be the set of automorphisms of $V_n(\mathcal{A})$ of the form

$$(E1.0.1) \quad g: \quad x_i \rightarrow r_i x_{\sigma(i)} \quad \forall i$$

where $\sigma \in S_n$ and $r_i \in k^\times = k \setminus \{0\}$. It is easy to see that g is an automorphism of $V_n(\mathcal{A})$ if and only if

$$(E1.0.2) \quad a_{ij} = r_i r_j a_{\sigma(i)\sigma(j)}$$

for all $i \neq j$. Such an automorphism g of $V_n(\mathcal{A})$ is denoted by $g(\sigma, \{r_1, \dots, r_n\})$. The following lemma is easy. We will use another obvious automorphism of $V_n(\mathcal{A})$, which sends x_i to $-x_i$ for all $i = 1, \dots, n$. We denote this automorphism by -1 , while the identity is just denoted by 1 .

Lemma 1.1. *Suppose that $n \geq 2$.*

- (1) [CPWZ1, Lemma 4.3] $G(\mathcal{A})$ is a group, and $\text{Aut}_{\text{af}}(V_n(\mathcal{A})) = G(\mathcal{A})$.
- (2) *There is a short exact sequence*

$$1 \rightarrow Z \rightarrow G(\mathcal{A}) \rightarrow S_n$$

where Z is a subgroup of $(k^\times)^n$. As a consequence, $G(\mathcal{A})$ is virtually abelian (so virtually solvable).

- (3) [CPWZ1, Theorem 4.9(3)] *If n is even, then $\text{Aut}(V_n(\mathcal{A})) = \text{Aut}_{\text{af}}(V_n(\mathcal{A})) = G(\mathcal{A})$.*

In parts (4–5) we further assume that $a_{ij} \neq 0$ for all $i < j$.

(4) If $n \geq 3$, there is a short exact sequence

$$1 \rightarrow \{\pm 1\} \rightarrow G(\mathcal{A}) \rightarrow S_n.$$

(5) If $n \geq 4$ is even, then $\text{Aut}(V_n(\mathcal{A})) = G(\mathcal{A})$ is finite.

Proof. (1) By a direct computation, $G(\mathcal{A}) \subseteq \text{Aut}_{\text{af}}(V_n(\mathcal{A}))$.

Now fix $g \in \text{Aut}_{\text{af}}(V_n(\mathcal{A}))$. By [CPWZ1, Lemma 4.3], there is a permutation $\sigma \in S_n$ and scalars $r_i \in k^\times$ such that $g(x_i) = r_i x_{\sigma(i)}$ for all i . Applying g to the relations of $V_n(\mathcal{A})$, one obtains that $a_{ij} = r_i r_j a_{\sigma(i)\sigma(j)}$ for all i, j . So $g \in G(\mathcal{A})$. Thus $\text{Aut}_{\text{af}}(V_n(\mathcal{A})) \subseteq G(\mathcal{A})$.

(2) The map from $G(\mathcal{A})$ to S_n takes $g(\sigma, \{r_1, \dots, r_n\})$ to σ . So the kernel Z of this map consists of all $g(\text{Id}, \{r_1, \dots, r_n\})$ where $a_{ij} = r_i r_j a_{ij}$ for all $i \neq j$. Since each r_i is in $k^\times$, Z is a subgroup of $(k^\times)^n$. Since Z is abelian, $G(\mathcal{A})$ is virtually abelian by definition.

(3) By [CPWZ1, Theorem 4.9(3)], $\text{Aut}(V_n(\mathcal{A})) = \text{Aut}_{\text{af}}(V_n(\mathcal{A}))$. The assertion follows from part (1).

(4) By the proof of part (2), Z consists of all $g(\text{Id}, \{r_1, \dots, r_n\})$ where $\{r_i\}_{i=1}^n$ satisfy $a_{ij} = r_i r_j a_{ij}$ for all $i \neq j$. Since $a_{ij} \neq 0$, we have $r_i r_j = 1$ for all $i < j$. Since $n \geq 3$, we obtain that $r_i = r_j$ for all $i < j$ and so $r_i = 1$ or $r_i = -1$.

(5) This follows from parts (3,4). $\square$

Let $G_1(\mathcal{A})$ denote the subgroup of $G(\mathcal{A})$ consisting of all $g(\sigma, \{r_1, \dots, r_n\})$ such that $\prod_{i=1}^n r_i = 1$. If $\mathcal{A} = \{1\}_{i < j}$ (i.e., the W_n case) and $n \geq 1$ is even, then $G(\mathcal{A}) = G_1(\mathcal{A})$ by [CPWZ1, Theorem 1]. We give an example for which $G(\mathcal{A}) \neq G_1(\mathcal{A})$ when $n = 4$.

Example 1.2. Let $n = 4$ and $(a_{ij})_{4 \times 4} = \begin{pmatrix} * & 1 & -1 & 1 \\ 1 & * & 1 & -a^2 \\ -1 & 1 & * & -a \\ 1 & -a^2 & -a & * \end{pmatrix}$ where $a^3 = -1$

and $a \neq -1$. It is easy to check that $g((123), \{1, 1, -1, a\})$ is in $G(\mathcal{A})$, but not in $G_1(\mathcal{A})$.

We refer to [JZ, Definition 0.6] for the definitions of Auslander-Gorenstein and GKdim-Macaulay.

Theorem 1.3. *Let G be a finite subgroup of $G_1(\mathcal{A})$. Suppose that $\text{char } k$ does not divide $|G|$. Then the fixed subring $V_n(\mathcal{A})^G$ is filtered Artin-Schelter Gorenstein, Auslander-Gorenstein and GKdim-Macaulay.*

Proof. First of all, we note that both $k_{-1}[x_1, \dots, x_n]$ and $V_n(\mathcal{A})$ are filtered Artin-Schelter Gorenstein, Auslander-Gorenstein and GKdim-Macaulay. We claim that, for each $g \in G \subset G_1(\mathcal{A})$, its homological determinant, denoted by $\text{hdet}_{V_n(\mathcal{A})} g$, is 1. If the claim holds, then the assertion follows from [JZ, Theorem 3.5].

Now we prove the claim. By definition [JZ, p. 677], $\text{hdet}_{V_n(\mathcal{A})} g = \text{hdet}_{\text{gr } V_n(\mathcal{A})} g$. To show the claim, we consider the algebra $\text{gr } V_n(\mathcal{A}) \cong k_{-1}[x_1, \dots, x_n] := B$ and identify G with a subgroup of $\text{Aut}(B)$ in the natural way. For any element $g = g(\sigma, \{r_1, \dots, r_n\}) \in G$, there is a decomposition

$$g(\sigma, \{r_1, \dots, r_n\}) = g(\sigma, \{1, \dots, 1\}) \circ g(\text{Id}, \{r_1, \dots, r_n\})$$

as in $\text{Aut}(B)$. Then the homological determinant of $g(\sigma, \{1, \dots, 1\})$ is 1 by the proof of [KKZ1, Theorem 1.5]. The homological determinant of $g(\text{Id}, \{r_1, \dots, r_n\})$

is $\prod_{i=1}^n r_i$, which is 1 by the definition of $G_1(\mathcal{A})$. Combining these, we see that $\text{hdet}_B g = 1$ for all $g \in G$. Thus we have proved the claim, and therefore the theorem. $\square$

Now we are ready to prove Theorem 1.

Theorem 1.4. *Let n be an even integer ≥ 4 and G a group acting on W_n faithfully. Suppose that $\text{char } k$ does not divide $2|G|$. Then the fixed subring W_n^G is filtered Artin-Schelter Gorenstein, Auslander-Gorenstein, and GKdim-Macaulay.*

Proof. Since n is even and $\text{char } k \neq 2$, by [CPWZ1, Theorem 1], $\text{Aut}(W_n) = S_n \times \{\pm 1\}$. So G is a subgroup of $S_n \times \{\pm 1\}$, which is $G_1(\{1\}_{i < j})$ when n is even. The assertion follows from Theorem 1.3. $\square$

Remark 1.5. (1) Theorem 1 is clearly a consequence of Theorem 1.4: if the group G acts on W_n , then $W_n^G = W_n^{\text{im } G}$, where $\text{im } G$ is the image of G in $\text{Aut } W_n$. Note that $\text{im } G$ acts faithfully.
(2) The hypothesis that “ $\text{char } k$ does not divide $2|G|$ ” is missing from [CPWZ1, Theorem 2]. We do not know whether W_n^G is filtered Artin-Schelter Gorenstein when $\text{char } k$ divides $2|G|$.
(3) By [CWWZ, Example 3.4(2) for D_4], the fixed subring $W_2^{S_2 \times \{\pm 1\}}$ is a commutative Gorenstein algebra with isolated singularities. So we expect that, in general, W_n^G is not regular. (Note that [CWWZ, Example 3.4(2)] was stated in terms of the ring $R = k\langle u, v \rangle / (u^2 + v^2 - 1)$, but this is isomorphic to W_2 .)

For completeness we have the following assertion for $n = 2$; see also [CWWZ, Theorem 0.1].

Proposition 1.6. *Let G be a finite group acting on W_2 faithfully. Suppose that $\text{char } k$ does not divide $2|G|$. Then the fixed subring W_2^G is filtered Artin-Schelter Gorenstein, Auslander-Gorenstein, and GKdim-Macaulay.*

Proof. By [AD], $\text{Aut}(W_2) = S_2 \ltimes k^\times$. So G is a finite subgroup of $S_2 \ltimes k^\times$. Note that the G -action preserves the given filtration of W_2 . By [CWWZ, Lemma 2.7], the homological determinant of the G -action is trivial (namely, $\text{hdet}_{W_2} g = 1$ for all $g \in G$). The assertion follows from [JZ, Theorem 3.5]. $\square$

By [KKZ1, Theorem 1.5(2)], if G is a subgroup of S_n other than $\{1\}$, then $k_{-1}[x_1, \dots, x_n]^G$ is not regular: its global dimension is infinite. When G is a special subgroup of S_n , then $k_{-1}[x_1, \dots, x_n]^G$ is a classical complete intersection [KKZ1, Theorem 3.12]. Some version of this should be true for W_n . Here we only give one simple example.

Example 1.7. Some of the arguments in this example were motivated by [KKZ1, Example 3.1].

Let S_2 act on W_2 in the obvious way. Then $A := W_2^{S_2}$ is a complete intersection, in the sense that it is isomorphic to $B/(f)$ for some regular algebra B and some normal nonzerodivisor $f \in B$. To see this, we note that A is generated by $u = x_1 + x_2$ and $v = x_1^3 + x_2^3$ satisfying two relations $[u^2, v] = [u, v^2] = 0$. By a straightforward computation, we have

$$r_3 := 2u^6 - 3u^3v - 3vu^3 + 4v^2 - 5u^4 + 3u^2 + 4(uv + vu) = 0.$$

One can check that A is isomorphic to $B/(r_3)$, where B is the down-up algebra $k\langle u, v \rangle / ([u^2, v] = [u, v^2])$.

Proposition 1.8. *Let G be the subgroup $\{\pm 1\}$ of $\text{Aut}(V_n(\mathcal{A}))$. If $a_{ij} \neq 0$ for some $i < j$, then $V_n(\mathcal{A})^G$ is regular.*

Proof. Define $\deg x_i = 1 \in \mathbb{Z}/(2)$. This makes $V_n(\mathcal{A})$ into a $\mathbb{Z}/(2)$ -graded algebra. Since the automorphism -1 will act on a length s monomial $m = x_{i_1} \dots x_{i_s}$ by multiplication by $(-1)^s$, such a monomial is in $V_n(\mathcal{A})^G$ if and only if s is even, which holds if and only if m is in the degree 0 part $(V_n(\mathcal{A}))_0$ of $V_n(\mathcal{A})$. Thus $V_n(\mathcal{A})^G = (V_n(\mathcal{A}))_0$. Since $x_i x_j + x_j x_i = a_{ij} \neq 0$ for some $i < j$, $V_n(\mathcal{A})$ is strongly $\mathbb{Z}/(2)$ -graded. It is well known that $V_n(\mathcal{A})$ is regular, so $(V_n(\mathcal{A}))_0$ is regular of the same global dimension by a theorem of Dade's [Da, Theorem 2.8] that the categories of $(V_n(\mathcal{A}))_0$ -modules and $\mathbb{Z}/(2)$ -graded $V_n(\mathcal{A})$ -modules are equivalent. $\square$

2. PROOF OF THEOREM 2

In this section we assume that k is a commutative domain.

Let A be an algebra generated over k by $x_1, x_2, \dots, x_t$ for some $t \geq 3$, and let R be the subalgebra generated by $x_3, \dots, x_t$. Suppose we have an algebra automorphism of A of the following form:

$$(E2.0.1) \quad g(x_i) = \begin{cases} x_1 + a_0 + a_1 x_2 & i = 1, \\ x_i & i \neq 1, \end{cases}$$

where a_0, a_1 are in the subalgebra R , and in particular are fixed by g . The following lemma is easy.

Lemma 2.1. *Let n be an integer. Then*

$$g^n(x_i) = \begin{cases} x_1 + na_0 + na_1 x_2 & i = 1, \\ x_i & i \neq 1. \end{cases}$$

Assume that, when n and m are any elements in k , the following are algebra isomorphisms of A :

$$(E2.1.1) \quad g^n(x_i) = \begin{cases} x_1 + na_0 + na_1 x_2 & i = 1, \\ x_i & i \neq 1, \end{cases}$$

and

$$(E2.1.2) \quad h^m(x_i) = \begin{cases} x_2 + mb_0 + mb_1 x_1 & i = 2, \\ x_i & i \neq 2. \end{cases}$$

This is automatic if $\text{char } k = 0$. We abuse notation and denote these by g^n and h^m , even if n and m are not integers. It is easy to check that $g^n g^{n'} = g^{n+n'}$.

By symmetry, if we also assume that we have an algebra automorphism

$$(E2.1.3) \quad h(x_i) = \begin{cases} x_2 + b_0 + b_1 x_1 & i = 2, \\ x_i & i \neq 2, \end{cases}$$

where $b_0, b_1 \in R$, then for any $m \in k$, we have

$$h^m(x_i) = \begin{cases} x_2 + mb_0 + mb_1 x_1 & i = 2, \\ x_i & i \neq 2. \end{cases}$$

Lemma 2.2. *For any $m, n \in k$,*

$$h^m g^n(x_i) = \begin{cases} x_1 + (na_0 + na_1 mb_0) + na_1 x_2 + na_1 mb_1 x_1 & i = 1, \\ x_2 + mb_0 + mb_1 x_1 & i = 2, \\ x_i & i \neq 1, 2, \end{cases}$$

and

$$g^n h^m(x_i) = \begin{cases} x_1 + na_0 + na_1 x_2 & i = 1, \\ x_2 + (mb_0 + mb_1 na_0) + mb_1 x_1 + mb_1 na_1 x_2 & i = 2, \\ x_i & i \neq 1, 2. \end{cases}$$

Let z be a positive integer and $n_1, \dots, n_z, m_1, \dots, m_z$ nonzero elements in k . Define automorphisms

$$\begin{aligned} \tau_z &= g^{n_z} h^{m_{z-1}} g^{n_{z-1}} \dots h^{m_1} g^{n_1}, \\ \sigma_z &= h^{m_z} \tau_z = h^{m_z} g^{n_z} h^{m_{z-1}} g^{n_{z-1}} \dots h^{m_1} g^{n_1}. \end{aligned}$$

Lemma 2.3. *Retain the above notation. We have*

$$\tau_z(x_1) = d_0 + d_1 x_1 + d_2 x_2 + \left(\prod_{s=1}^{z-1} m_s \right) \left(\prod_{s=1}^z n_s \right) a_1 (b_1 a_1)^{z-1} x_2$$

where d_0, d_1, d_2 are elements in R . Further,

$$d_1 = \sum_{s=0}^{z-1} \alpha_s (a_1 b_1)^s \quad \text{and} \quad d_2 = \sum_{s=0}^{z-2} \beta_s a_1 (b_1 a_1)^s$$

where α_s, β_s are elements in the subring of k generated by $n_1, \dots, n_z, m_1, \dots, m_z$.

Proof. When $z = 1$, this is Lemma 2.1. Now assume that $z > 1$. By definition, $\tau_z = g^{n_z} h^{m_{z-1}} \tau_{z-1}$. By a direct computation and the induction hypothesis,

$$\begin{aligned} \tau_z(x_1) &= g^{n_z} h^{m_{z-1}} \tau_{z-1}(x_1) \\ &= g^{n_z} h^{m_{z-1}} (d'_0 + d'_1 x_1 + d'_2 x_2 + \left(\prod_{s=1}^{z-2} m_s \right) \left(\prod_{s=1}^{z-1} n_s \right) a_1 (b_1 a_1)^{z-2} x_2) \\ &= d'_0 + d'_1 (x_1 + n_z a_0 + n_z a_1 x_2) \\ &\quad + d'_2 (x_2 + (m_{z-1} b_0 + m_{z-1} b_1 n_z a_0) + m_{z-1} b_1 x_1 + m_{z-1} b_1 n_z a_1 x_2) \\ &\quad + \left(\prod_{s=1}^{z-2} m_s \right) \left(\prod_{s=1}^{z-1} n_s \right) a_1 (b_1 a_1)^{z-2} \\ &\quad \quad (x_2 + (m_{z-1} b_0 + m_{z-1} b_1 n_z a_0) + m_{z-1} b_1 x_1 + m_{z-1} b_1 n_z a_1 x_2) \\ &= d_0 + d_1 x_1 + d_2 x_2 + \left(\prod_{s=1}^{z-1} m_s \right) \left(\prod_{s=1}^z n_s \right) a_1 (b_1 a_1)^{z-1} x_2 \end{aligned}$$

where

$$\begin{aligned}
d_0 &= d'_0 + d'_1 n_z a_0 + d'_2 (m_{z-1} b_0 + m_{z-1} b_1 n_z a_0) \\
&\quad + \left(\prod_{s=1}^{z-2} m_s \right) \left(\prod_{s=1}^{z-1} n_s \right) a_1 (b_1 a_1)^{z-2} (m_{z-1} b_0 + m_{z-1} b_1 n_z a_0), \\
d_1 &= d'_1 + d'_2 m_{z-1} b_1 + \left(\prod_{s=1}^{z-2} m_s \right) \left(\prod_{s=1}^{z-1} n_s \right) a_1 (b_1 a_1)^{z-2} m_{z-1} b_1, \\
d_2 &= d'_1 n_z a_1 + d'_2 + d'_2 m_{z-1} b_1 n_z a_1 + \left(\prod_{s=1}^{z-2} m_s \right) \left(\prod_{s=1}^{z-1} n_s \right) a_1 (b_1 a_1)^{z-2}.
\end{aligned}$$

The formulas for d_1 and d_2 follow easily by induction. $\square$

In the next proposition, the free product $G * H$ of two groups G and H will be used. It is well known that every element in $G * H$ can be expressed uniquely in the form $g_1 h_1 g_2 h_2 \cdots g_s h_s$ for some s and for some $g_i \in G$, $h_i \in H$ with $h_i \neq 1$ for $i \neq s$ and $g_i \neq 1$ for $i \neq 1$. The following lemma is well known.

Lemma 2.4. *Let G be a group of order at least 3. Then $G * G$ contains a free subgroup of rank two.*

Proof. If $x \in G$ has infinite order, then $\langle x \rangle * \langle x \rangle$ is a free group of rank two contained in $G * G$.

If x has order $n \geq 3$, then $\langle x \rangle * \langle x \rangle$ contains a free subgroup of rank $(n-1)^2$ [AP, Theorem 1].

If every element x in G has order 2, then G is abelian and G contains $(\mathbb{Z}/(2))^{\oplus 2}$. It is well known that $(\mathbb{Z}/(2))^{\oplus 2} * (\mathbb{Z}/(2))^{\oplus 2}$ contains a free subgroup of rank $(4-1)^2$ [AP, Theorem 1]. $\square$

Let k_0 be a subring of k . We say an element $f \in A$ is *transcendental over k_0* if there are no $r_i \in k_0$ with $r_n \neq 0$ such that $\sum_{i=0}^n r_i f^i = 0$.

Proposition 2.5. *Retain the hypotheses and notation as above; in particular, assume that g^n and h^m from (E2.1.1) and (E2.1.2) are automorphisms of A for all $n, m \in k$. Assume that $R + Rx_1 + Rx_2$ is a free left R -module with basis $\{1, x_1, x_2\}$ and that the element $a_1 b_1 \in A$ is transcendental over a subring $k_0 \subseteq k$.*

- (1) *Suppose m_s and n_s are nonzero elements in k_0 for all s . Then neither σ_z nor τ_z is the identity automorphism.*
- (2) *$\text{Aut}(A)$ contains the free product $(k_0, +) * (k_0, +)$. As a consequence, if $k_0 \neq \mathbb{Z}/(2)$, then $\text{Aut}(A)$ contains a free subgroup of rank two.*

Proof. (1) Since $a_1 b_1$ is transcendental over k_0 , $d_2 + (\prod_{s=1}^{z-1} m_s) (\prod_{s=1}^z n_s) a_1 (b_1 a_1)^{z-1}$ is not zero. Since $R + Rx_1 + Rx_2$ is free, $\tau_z(x_1) \neq x_1$ by Lemma 2.3. So τ_z is not the identity. Since $h^{-m_z}(x_1) = x_1$, we deduce that $\tau_z \neq h^{-m_z}$, so σ_z is not the identity.

(2) There is a natural group map $(k_0, +) * (k_0, +) \rightarrow \text{Aut}(A)$ sending $m_z * n_z * m_{z-1} * n_{z-1} * \cdots * m_1 * n_1$ to $h^{m_z} g^{n_z} h^{m_{z-1}} g^{n_{z-1}} \cdots h^{m_1} g^{n_1}$. By part (1) the kernel of this map is $\{1\}$. So the main assertion follows. The consequence follows from Lemma 2.4. $\square$

Here is a version of [CPWZ1, Lemma 5.11] for the algebra $V_n(\mathcal{A})$. First recall from [CPWZ1, Example 5.10(2)],

$$\Omega(x_1, \dots, x_t) := \sum_{\sigma \in S_t} (-1)^{|\sigma|} x_{\sigma(1)} \dots x_{\sigma(t)}.$$

Lemma 2.6. *For any $t \leq n$, consider the element $\Omega_t = \Omega(x_1, \dots, x_t)$ in $V_n(\mathcal{A})$. Then $x_i \Omega_t = (-1)^{n-1} \Omega_t x_i$ for all $i = 1, \dots, t$.*

Proof. The proof is similar to that of [CPWZ1, Lemma 5.11] and is therefore omitted. $\square$

Now we are ready to prove Theorem 2.

Proof of Theorem 2. Let $n \geq 3$ be odd. By Lemma 2.6, we have two automorphisms of $V_n(\mathcal{A})$, as in [CPWZ1, Example 5.10(2)]:

$$g : x_i \rightarrow \begin{cases} x_1 + \Omega(x_2, x_3, \dots, x_n) & i = 1, \\ x_i & 2 \leq i \leq n, \end{cases}$$

and

$$h : x_i \rightarrow \begin{cases} x_2 + \Omega(x_1, x_3, \dots, x_n) & i = 2, \\ x_i & i = 1 \text{ or } 3 \leq i \leq n. \end{cases}$$

Using the notation in (E2.0.1) and (E2.1.3), both a_1 and b_1 have leading term equal to $(-1)^{n-2}(n-1)! \prod_{i=3}^n x_i$. It is easy to see that $R + Rx_1 + Rx_2$ is a free left R -module with basis $\{1, x_1, x_2\}$, and it is clear that $a_1 b_1$ is transcendental over k . Since $k_0 = k \neq \mathbb{Z}/(2)$, the assertion follows from Proposition 2.5(2). $\square$

Remark 2.7. It was claimed in [CPWZ1, Example 5.10] that $\text{Aut}(W_n)$ is not affine when n is odd. This is true when $\text{char } k$ does not divide $(n-1)!$; it is not known what happens when $\text{char } k$ divides $(n-1)!$. The hypothesis that “ $\text{char } k$ does not divide $(n-1)!$ ” was missing from [CPWZ1, Example 5.10].

We conclude with a corollary of Proposition 2.5 which deals with skew polynomial rings. See [CPWZ3] for more results along these lines. Let $\{p_{ij} \in k^\times : 1 \leq i < j \leq n\}$ be a set of parameters. The *skew polynomial ring*, denoted by $k_{p_{ij}}[x_1, \dots, x_n]$, is defined to be the algebra generated by $x_1, \dots, x_n$ subject to the relations $x_j x_i = p_{ij} x_i x_j$ for all $i < j$. We set $p_{ji} = p_{ij}^{-1}$ and $p_{ii} = p_{jj} = 1$ for all $i < j$. For any $1 \leq s \leq n$, let

$$T_s := \left\{ (d_1, \dots, \widehat{d_s}, \dots, d_n) \in \mathbb{N}^{n-1} : \prod_{\substack{j=1 \\ j \neq s}}^n p_{ij}^{d_j} = p_{is} \quad \forall i \neq s \right\}.$$

In [CPWZ2, Theorem 3.8], it was proved that if each p_{ij} is a root of unity and $T_s = \emptyset$ for all s , then every automorphism of $k_{p_{ij}}[x_1, \dots, x_n]$ is affine. Note also that if each p_{ij} is a root of unity and some T_s is non-empty, then T_s is in fact infinite. If we drop the assumption on the p_{ij} 's and we allow at most one T_s to be infinite, then we can still understand the automorphism group of $k_{p_{ij}}[x_1, \dots, x_n]$ – see [CPWZ2, Theorem 4]. The following result concerns $\text{Aut}(k_{p_{ij}}[x_1, \dots, x_n])$ in another case.

Corollary 2.8. *Let $n \geq 3$ be an integer and let $A := k_{p_{ij}}[x_1, \dots, x_n]$ be a skew polynomial ring. Suppose that T_1 contains an element of the form $(1, d_3, \dots, d_n)$ and T_2 contains an element of the form $(1, d'_3, \dots, d'_n)$. Suppose that at least one of $d_3, \dots, d_n, d'_3, \dots, d'_n$ is nonzero. Then $\text{Aut}(A)$ contains $(k, +) * (k, +)$. If, further, $k \neq \mathbb{Z}/(2)$, then $\text{Aut}(A)$ contains a free group on two generators.*

Proof. For $d = (1, d_3, \dots, d_n) \in T_1$ and $d' = (1, d'_3, \dots, d'_n) \in T_2$ as in the statement, define $f_d := x_2 x_3^{d_3} \dots x_n^{d_n}$, $f_{d'} := x_1 x_3^{d'_3} \dots x_n^{d'_n}$. The maps, defined in [CPWZ2, (2.10.1)],

$$g(f_d, 1) : x_i \mapsto \begin{cases} x_i & \text{if } i \neq 1, \\ x_1 + f_d & \text{if } i = 1, \end{cases} \quad g(f_{d'}, 2) : x_i \mapsto \begin{cases} x_i & \text{if } i \neq 2, \\ x_2 + f_{d'} & \text{if } i = 2, \end{cases}$$

extend to algebra automorphisms of A . Using the notation (E2.0.1) and (E2.1.3), $a_1 = \prod_{j=3}^n p_{j,2}^{d_j} x_3^{d_3} \dots x_n^{d_n}$ and $b_1 = \prod_{j=3}^n p_{j,1}^{d'_j} x_3^{d'_3} \dots x_n^{d'_n}$. It is clear that $R + Rx_1 + Rx_2$ is a free left R -module with basis $\{1, x_1, x_2\}$ and that $a_1 b_1$ is transcendental over k . The assertion follows from Proposition 2.5(2). $\square$

Note that this applies to the ordinary polynomial ring: if $p_{ij} = 1$ for all i, j , then each set T_s is all of $\mathbb{N}^{n-1}$.

3. PROOF OF THEOREM 3

In this short section we assume that k is a commutative domain. The main idea is to use the discriminant to solve isomorphism questions. Given two algebras with similar properties, it is generally very difficult to determine whether they are isomorphic. When the discriminant controls the the automorphism groups of these algebras, it also controls the isomorphisms between them.

Given an algebra A , we write $d(A/Z(A))$ for the discriminant of A over its center, as defined in [CPWZ1] or [CPWZ2]. The following lemma is clear.

Lemma 3.1. *Suppose A and B are PI algebras, and let $g : A \rightarrow B$ be an algebra isomorphism. Then $g(d(A/Z(A))) = c d(B/Z(B))$ for some unit $c \in Z(B)$.*

We keep the notation for $V_n(\mathcal{A})$ and $V_n(\mathcal{A}')$ as above, except that it is convenient to distinguish between their generators, so we use $\{x_i\}$ for the generators of $V_n(\mathcal{A})$ and $\{x'_i\}$ for the generators of $V_n(\mathcal{A}')$.

Proof of Theorem 3. First of all, if there exist $\sigma \in S_n$ and invertible scalars λ_i such that $a'_{ij} = \lambda_i \lambda_j a_{\sigma(i)\sigma(j)}$, then the map g determined by $x'_i \mapsto \lambda_i x_{\sigma(i)}$ for $i = 1, \dots, n$ extends to an algebra isomorphism $V_n(\mathcal{A}') \rightarrow V_n(\mathcal{A})$.

Conversely, let $g : V_n(\mathcal{A}') \rightarrow V_n(\mathcal{A})$ be an algebra isomorphism. By [CPWZ1, Theorem 4.9],

$$d := d(V_n(\mathcal{A})/Z(V_n(\mathcal{A}))) = \left(\prod_{i=1}^n x_i^2 \right)^{2^{n-1}} + (\text{cwlt}),$$

where (cwlt) (“component-wise less than”) is a linear combination of terms with degree strictly less than the degree of $(\prod_{i=1}^n x_i^2)^{2^{n-1}}$. By symmetry,

$$d' := d(V_n(\mathcal{A}')/Z(V_n(\mathcal{A}'))) = \left(\prod_{i=1}^n (x'_i)^2 \right)^{2^{n-1}} + (\text{cwlt}).$$

By Lemma 3.1, $g(d') = cd$ for some unit $c \in Z(V_n(\mathcal{A})) = k[x_1^2, \dots, x_n^2]$. So c is a unit in $k^\times$. The equation $g(d') = cd$ implies that

$$(E3.1.1) \quad \left(\prod_{i=1}^n (g(x'_i))^2 \right)^{2^{n-1}} + g((\text{cwt})) = c \left(\prod_{i=1}^n x_i^2 \right)^{2^{n-1}} + (\text{cwt}).$$

Note that $g(x'_i)$ has degree at least 1 since it is not in k . So each term in $g((\text{cwt}))$ has degree strictly less than the degree of $(\prod_{i=1}^n (g(x'_i))^2)^{2^{n-1}}$. This implies that the degree of the left-hand term in (E3.1.1) is equal to the degree of $(\prod_{i=1}^n (g(x'_i))^2)^{2^{n-1}}$. The degree of the right-hand term of (E3.1.1) is $n2^n$. Therefore $\deg g(x'_i) = 1$ for all i . Thus if we use the standard filtration on both $V_n(\mathcal{A}')$ and $V_n(\mathcal{A})$, then g preserves the filtration. Consider the graded algebra isomorphism $\text{gr } g : \text{gr } V_n(\mathcal{A}') \rightarrow \text{gr } V_n(\mathcal{A})$, recalling that $\text{gr } V_n(\mathcal{A}) \cong k_{-1}[x_1, \dots, x_n]$. By [CPWZ1, Lemma 4.3], there are units $r_i \in k$ and a permutation $\sigma \in S_n$ such that $\text{gr } g(x'_i) = r_i x_{\sigma(i)}$. Going back to g , there must be $a_i \in k$ such that $g(x'_i) = r_i x_{\sigma(i)} + a_i$ for all i . Using the second half of the proof of [CPWZ1, Lemma 4.3], one sees that $a_i = 0$ for all i . The assertion follows. $\square$

4. EXAMPLES AND QUESTIONS

In this final section, we give some examples and list some questions concerning the algebra $V_n(\mathcal{A})$. For the sake of convenience, we assume that k is an algebraically closed field of characteristic zero.

Lemma 4.1. *If the order of $\text{Aut}(V_n(\mathcal{A}))$ is $2n!$, then n is an even integer ≥ 4 and $V_n(\mathcal{A})$ is isomorphic to W_n . As a consequence, $\text{Aut}(V_n(\mathcal{A})) \cong S_n \times \{\pm 1\}$.*

Proof. By Theorem 2, n is even. If $a_{ij} = 0$ for all $i < j$, then $\text{Aut}(V_n(\mathcal{A})) \cong S_n \ltimes (k^\times)^n$ which does not have order $2n!$. So $a_{ij} \neq 0$ for some $i < j$.

If $n = 2$, then a_{12} must be nonzero, in which case $V_2(\mathcal{A})$ is isomorphic to W_2 . By [AD], $\text{Aut}(W_2) = S_2 \ltimes k^\times$, so $n \neq 2$. Hence n is an even integer larger than or equal to 4.

Given that $a_{ij} \neq 0$ for some $i < j$, we claim that $a_{st} \neq 0$ for all $s < t$. By Lemma 1.1, $|\text{Aut}(V_n(\mathcal{A}))| = 2m$, where m is the order of the image of $\text{Aut}(V_n(\mathcal{A}))$ in S_n . Since $|\text{Aut}(V_n(\mathcal{A}))|$ is $2n!$, the map $\text{Aut}(V_n(\mathcal{A})) \rightarrow S_n$ must be surjective, so there is a short exact sequence

$$1 \rightarrow \{\pm 1\} \rightarrow \text{Aut}(V_n(\mathcal{A})) \rightarrow S_n \rightarrow 1.$$

This means that for each $\sigma \in S_n$, there is a corresponding element $g(\sigma, \{r_i\}) \in \text{Aut}(V_n(\mathcal{A}))$. In particular, given s and t with $1 \leq s < t \leq n$, there is a permutation $\sigma \in S_n$ such that $(\sigma(i), \sigma(j)) = (s, t)$. By (E1.0.2), $a_{ij} = r_i r_j a_{\sigma(i)\sigma(j)} = r_i r_j a_{st}$. Since $a_{ij} \neq 0$, we obtain that $a_{st} \neq 0$.

Since elements in k have square roots in k , after replacing x_i by $b_i x_i$ for $i = 1, 2, 3$ for appropriate b_1, b_2, b_3 , we may assume that $a_{12} = a_{13} = a_{23} = 1$. For each $i > 3$, by replacing x_i by $a_{1i}^{-1} x_i$, we may assume that $a_{1i} = 1$ for all $i > 3$. Summarizing, we have $a_{1i} = 1$ for all $i > 1$ and $a_{23} = 1$.

Next we claim that $a_{ij} = 1$ for all $i < j$. First we show that $a_{2i} = 1$ for all $i \neq 2$. Taking an element $g((12), \{r_i\})$ in $\text{Aut}(V_n(\mathcal{A}))$, (E1.0.2) implies that

$$\begin{aligned} a_{12} &= r_1 r_2 a_{21}, \\ a_{13} &= r_1 r_3 a_{23}, \\ a_{23} &= r_2 r_3 a_{13}. \end{aligned}$$

Since $a_{12} = a_{13} = a_{23} = 1$, $r_1 = r_2 = r_3 = \pm 1$. By (E1.0.2),

$$a_{3i} = r_3 r_i a_{3i} \quad \forall i > 3.$$

Hence $r_i = r_3 = \pm 1$ for all $i > 3$. So $r_i = r_1 = \pm 1$ for all i . By (E1.0.2) again, $a_{2i} = r_2 r_i a_{1i} = 1$ for all i . By symmetry, $a_{3i} = 1$ for all $i \neq 3$. Secondly, we show that $a_{ij} = 1$ for all $3 < i < j$. Nothing needs to be proved if $n = 4$. For $n \geq 6$, it suffices to show $a_{45} = 1$ by symmetry. Taking $g((14), \{r_i\})$ in $\text{Aut}(V_n(\mathcal{A}))$, (E1.0.2) implies that

$$\begin{aligned} 1 &= a_{12} = r_1 r_2 a_{42} = r_1 r_2, \\ 1 &= a_{13} = r_1 r_3 a_{43} = r_1 r_3, \\ 1 &= a_{23} = r_2 r_3 a_{23} = r_2 r_3, \\ 1 &= a_{34} = r_3 r_4 a_{31} = r_3 r_4. \end{aligned}$$

Then $r_1 = r_2 = r_3 = r_4 = \pm 1$. By (E1.0.2),

$$a_{3i} = r_3 r_i a_{3i}, \quad \forall i > 4.$$

Hence $r_i = r_3 = \pm 1$ for all $i > 4$. So $r_i = r_1 = \pm 1$ for all i . By (E1.0.2) again, $a_{45} = r_4 r_5 a_{15} = 1$, as desired.

The consequence follows from [CPWZ1, Theorem 1]. $\square$

There are cases when $\text{Aut}(V_n(\mathcal{A}))$ is smaller than $S_n \times \{\pm 1\}$, as the next example shows.

Example 4.2. Let $n = 4$.

(1) Let q be transcendental over $\mathbb{Q} \subseteq k$. Let $a_{12} = q$, $a_{13} = q^2$, $a_{14} = q^4$, $a_{23} = q^8$, $a_{24} = q^{16}$, and $a_{34} = q^{32}$. For any $\{i_1, \dots, i_4\} = \{j_1, \dots, j_4\} = \{1, \dots, 4\}$, unless $i_s = j_s$ for all s , the element $a_{i_1 i_2} a_{i_3 i_4} a_{j_1 j_2}^{-1} a_{j_3 j_4}^{-1}$ is a non-trivial power of q , which is not a root of unity. Using (E1.0.1), (E1.0.2) and the fact that the homological determinant of g is $r_1 r_2 r_3 r_4$, one can show that

$$G(\mathcal{A})/\{\pm 1\} \cong \{Id, (12)(34), (13)(24), (14)(23)\} \cong (\mathbb{Z}/(2))^{\oplus 2}.$$

In general, one can show that when $n = 4$ and $a_{ij} \neq 0$ for all $i < j$, then $G(\mathcal{A})/\{\pm 1\}$ always contains the subgroup $\{Id, (12)(34), (13)(24), (14)(23)\}$.

(2) Let q be transcendental over $\mathbb{Q} \subseteq k$. Let $a_{12} = q$, $a_{13} = q^2$, $a_{14} = q^4$, $a_{23} = q^8$, and $a_{24} = a_{34} = 0$. We claim that $\text{Aut}(V_n(\mathcal{A})) \cong S_2 \times \{\pm 1\}$. If $g = g(\sigma, \{r_i\})$ is in $\text{Aut}(V_n(\mathcal{A}))$, then σ fixes 1 and 4, as 1 is the only index so that $a_{1i} \neq 0$ for all i and 4 is the only index so that $a_{i4} = 0$ for two different i . If g is neither 1 nor -1 , g must be of the form $g((23), \{r_i\})$. In fact, one can check that $G(\mathcal{A})/\{\pm 1\} \cong \{(23)\} \cong S_2$.

(3) Let $(a_{ij})_{4 \times 4} = \begin{pmatrix} * & 1 & -1 & 1 \\ 1 & * & 1 & -a^2 \\ -1 & 1 & * & -a \\ 1 & -a^2 & -a & * \end{pmatrix}$ where $a^3 = -1$ and $a \neq -1$. This is

the same as Example 1.2. One can show that $G(\mathcal{A})/\{\pm 1\} \cong A_4$.

Example 4.3. Let $n = 6$.

(1) Suppose that $a_{15} = a_{45} = a_{i6} = 0$ for all $i = 2, 3, 4, 5$, that all other a_{ij} are nonzero, and that $a_{12}a_{34} \neq a_{13}a_{24}$. Then $\text{Aut}(V_n(\mathcal{A})) \cong \{\pm 1\}$. To see this, we first note that 6 is the only index so that there are 4 different i such that $a_{i6} = 0$; thus if $g(\sigma, \{r_i\})$ is in $\text{Aut}(V_n(\mathcal{A}))$, σ fixes 6. Similarly, σ fixes 1, 4 and 5. The only possible nontrivial σ is (23). In the case of $g((23), \{r_i\})$, we have

$$\begin{aligned} a_{12} &= r_1 r_2 a_{13}, \\ a_{13} &= r_1 r_3 a_{12}, \\ a_{23} &= r_2 r_3 a_{23}, \\ a_{14} &= r_1 r_4 a_{14}, \\ a_{24} &= r_2 r_4 a_{34}. \end{aligned}$$

By the first three equations, we have $r_1^2 = 1$ and $r_2 r_3 = 1$. So $r_1 = \pm 1$. By the fourth equation, $r_4 = r_1$. Then the first and fifth equations contradict the hypothesis that $a_{12}a_{34} \neq a_{13}a_{24}$. Therefore the only $g \in \text{Aut}(V_n(\mathcal{A}))$ is of the form ± 1 , so $\text{Aut}(V_n(\mathcal{A})) \cong \{\pm 1\}$.

(2) Suppose that $a_{12} = 0$ and that $a_{ij} = 1$ for all $i < j$ except for a_{12} . Then $\text{Aut}(V_n(\mathcal{A}))/\{\pm 1\} \cong S_2 \times S_4$ and therefore $\text{Aut}(V_n(\mathcal{A})) \cong (S_2 \times S_4) \times \{\pm 1\}$.

Question 4.4. Which finite groups can be realized as $\text{Aut}(V_n(\mathcal{A}))$? For which of those groups G can one classify the algebras $V_n(\mathcal{A})$ such that $\text{Aut}(V_n(\mathcal{A})) \cong G$?

Similar to Theorem 1, we have the following. Recall that $G_1(\mathcal{A})$ is defined just after Lemma 1.1.

Lemma 4.5. *Suppose there is a short exact sequence*

$$1 \rightarrow \{\pm 1\} \rightarrow \text{Aut}(V_n(\mathcal{A})) \rightarrow H \rightarrow 1$$

where H is a subgroup of S_n such that $H = [H, H]$ (for example, $H = A_n$ when $n \geq 6$). Then

- (1) $G(\mathcal{A}) = G_1(\mathcal{A})$.
- (2) For each subgroup $G \subseteq \text{Aut}(V_n(\mathcal{A}))$, the fixed subring $V_n(\mathcal{A})^G$ is filtered AS Gorenstein.

Proof. Part (2) is a consequence of part (1) and Theorem 1.3, so we only prove part (1). Since n must be even, the homological determinant $\text{hdet} : \text{Aut}(V_n(\mathcal{A})) \rightarrow k^\times$ maps ± 1 to 1. Hence it induces a group homomorphism $\text{hdet}' : H \rightarrow k^\times$. Since $H = [H, H]$ and $k^\times$ is abelian, hdet' is the trivial map. So the image of hdet is $\{1\}$. This is equivalent to saying that $G(\mathcal{A}) = G_1(\mathcal{A})$. $\square$

Related to the above and Theorem 1, we have the following question.

Question 4.6. Classify all $V_n(\mathcal{A})$ (when n is even) such that $V_n(\mathcal{A})^G$ is Gorenstein for all subgroups $G \subseteq \text{Aut}(V_n(\mathcal{A}))$.

Theorem 3 suggests the following question.

Question 4.7. When n is odd, determine when two $V_n(\mathcal{A})$ s are isomorphic.

A related question is the following.

Question 4.8. If $V_n(\mathcal{A})$ is Morita equivalent to $V_n(\mathcal{A}')$, is $V_n(\mathcal{A})$ isomorphic to $V_n(\mathcal{A}')$?

Some Hopf algebra actions on W_2 are given in [CWWZ, Example 3.4]. It would be very interesting to work out all possible Hopf algebra actions.

Question 4.9. Suppose n is even. Classify all finite dimensional Hopf algebras that can act on W_n inner-faithfully.

Acknowledgments. S. Ceken was supported by the Scientific and Technological Research Council of Turkey (TUBITAK), Science Fellowships and Grant Programmes Department (Programme no. 2214). Y.H. Wang was supported by the Natural Science Foundation of China (grant nos. 10901098 and 11271239), the Scientific Research Starting Foundation for the Returned Overseas Chinese Scholars, Ministry of Education of China and the Innovation program of Shanghai Municipal Education Commission. J. J. Zhang was supported by the US National Science Foundation (Nos. DMS-0855743 and DMS-1402863).

REFERENCES

- [AD] J. Alev and F. Dumas, Rigidité des plongements des quotients primitifs minimaux de $U_q(sl(2))$ dans l'algèbre quantique de Weyl-Hayashi, Nagoya Math. J. **143** (1996), 119–146.
- [AP] M. Anshel and R. Prener, On free products of finite abelian groups. Proc. Amer. Math. Soc. **34** (1972), 343–345.
- [BJ] V.V. Bavula and D.A. Jordan, Isomorphism problems and groups of automorphisms for generalized Weyl algebras. Trans. Amer. Math. Soc. **353** (2001), no. 2, 769–794.
- [CPWZ1] S. Ceken, J. Palmieri, Y.-H. Wang and J.J. Zhang, The discriminant controls automorphism groups of noncommutative algebras, Adv. Math., **269** (2015), 551–584.
- [CPWZ2] S. Ceken, J. Palmieri, Y.-H. Wang and J.J. Zhang, The discriminant criterion and automorphism groups of quantized algebras, preprint (2014), arXiv:1402.6625
- [CPWZ3] S. Ceken, J. Palmieri, Y.-H. Wang and J.J. Zhang, Tits alternative for automorphisms and derivations, in preparation (2015).
- [CWWZ] K. Chan, C. Walton, Y.-H. Wang and J.J. Zhang, Hopf actions on filtered regular algebras, J. Algebra, **397** (2014), 68–90.
- [Ch] C. Chevalley, Invariants of finite groups generated by reflections, Amer. J. Math. **77** (1955), 778–782.
- [Da] E. Dade, Group-graded rings and modules, Math. Z. **174** (1980), 241–262.
- [JZ] N. Jing and J.J. Zhang, Gorensteinness of invariant subrings of quantum algebras, J. Algebra **221** (1999), no. 2, 669–691.
- [Ki] E. Kirkman, Invariant theory of Artin-Schelter regular algebras: a survey, preprint, 2014.
- [KKZ1] E. Kirkman, J. Kuzmanovich and J.J. Zhang, Invariants of (-1) -skew polynomial rings under permutation representations, Contemporary Mathematics, **623** (2014), 155–192.
- [KKZ2] E. Kirkman, J. Kuzmanovich and J.J. Zhang, Invariant theory of finite group actions on down-up algebras, Transform. Groups, DOI: 10.1007/S00031-014-9279-4.
- [ST] G. C. Shephard and J. A. Todd, Finite unitary reflection groups, Canadian J. Math. **6** (1954), 274–304
- [SAV] M. Suárez-Alvarez and Q. Vivas, Automorphisms and isomorphism of quantum generalized Weyl algebras, preprint (2012), arXiv:1206.4417v1.
- [Ti] J. Tits, Free subgroups in linear groups, J. Algebra **20** (1972), 250–270.
- [Wa] K. Watanabe, Certain invariant subrings are Gorenstein I, II, Osaka J. Math. **11** (1974) 1–8, 379–388.
- [Y1] M. Yakimov, The Andruskiewitsch-Dumas conjecture, Selecta Math. (N.S.) **20** (2014), no. 2, 421–464.
- [Y2] M. Yakimov, The Launois-Lenagan conjecture, J. Algebra **392** (2013), 1–9.

CEKEN: DEPARTMENT OF MATHEMATICS, AKDENİZ UNIVERSITY, 07058 ANTALYA, TURKEY
E-mail address: cekensecil@gmail.com

PALMIERI: DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98195, USA
E-mail address: palmieri@math.washington.edu

WANG: SCHOOL OF MATHEMATICS, SHANGHAI UNIVERSITY OF FINANCE AND ECONOMICS, SHANGHAI 200433, CHINA

E-mail address: yhw@mail.shufe.edu.cn

ZHANG: DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98195, USA

E-mail address: zhang@math.washington.edu